

# Outcomes of Neonates Requiring Surgical Interventions in Eldoret

Kasede M Ikol,<sup>1</sup> Peter W Saula,<sup>1</sup> Peter Gisore,<sup>1</sup> Everlyne Mvungu,<sup>2</sup> Henry R Mwangi<sup>2</sup>

<sup>1</sup> School of Medicine, Moi University

<sup>2</sup> Moi Teaching and Referral Hospital

**Correspondence to:** Dr. Peter Saula, PO Box 1998 - 30100 Eldoret; email: [saulapw@yahoo.com](mailto:saulapw@yahoo.com)

## Abstract

**Background:** Neonatal surgical conditions account for 11% of the Global Burden of Diseases (GBD); neglect of their management has been reported though most are amenable to surgery. Timely surgical interventions play a major role in determining better outcomes, thus improving quality of life and reducing disability. Several factors—sepsis control, care in newborn intensive care unit and availability of total parenteral nutrition—have been shown to improve the outcomes of neonates with surgical conditions. **Objective:** To evaluate the outcomes of neonates with conditions requiring surgical interventions. **Methods:** A prospective descriptive study in the Newborn Unit, Moi Teaching and Referral Hospital (MTRH). Data on the following study variables were obtained and analyzed: maternal age, place and mode of delivery, newborn's age at admission, birth weight, surgical condition, co-morbid conditions, treatment outcomes (discharge, death or referral to Kenyatta National Hospital for specialized care), surgical complications, time-to-initiation of oral feeds post-operatively, antenatal history and laboratory parameters; and length of hospital stay. **Results:** A total of 124 neonates were recruited; male to female ratio was 1.1:1 and median age at admission was 2 days (IQR 1, 5). Most (59.7%) were in the birth weight

range of 2.5–3.9 kg. The leading neonatal surgical conditions were gastroschisis 33(26.6%), neural tube defects 25(20.2%) and ARM 25(20.2%). Overall mortality rate was 31.5%. Antenatal clinic attendance was associated with decreased odds of mortality (OR 0.126; 95% CI 0.025–0.6429;  $p=0.013$ ), while neonatal sepsis, respiratory distress and electrolyte imbalance were associated with increased odds of mortality ([OR 3.4; 95% CI 1.09–22.06;  $p=0.049$ ], [OR 4.9; CI 0.91–11.61;  $p=0.001$ ] and [OR 3.1; CI 1.21–31.60;  $p=0.029$ ], respectively). **Conclusions:** The overall mortality rate of neonates requiring surgical interventions at MTRH was 31.5% and the median length of hospital stay was 14.5 days. Co-morbid conditions that increased the odds of mortality among these neonates were neonatal sepsis, respiratory distress and electrolyte imbalance.

**Key words:** Neonates, Surgical, Interventions, Outcomes  
Ann Afr Surg. 2019; 16 (1):20–25

DOI:<http://dx.doi.org/10.4314/aas.v16i1.5>

**Conflicts of Interest:** None

**Funding:** None

© 2019 Author. This work is licensed under the Creative Commons Attribution 4.0 International License.

## Introduction

The World Health Organization defines a neonate as child less than 28 days of age, a critical period during which the child is at the highest risk of dying (1). Neonatal surgical conditions account for 11% of the Global Burden of Diseases (GBD), most of which are congenital anomalies (2). Despite this, neglect is reported in their management, yet most of these conditions are amenable to surgery (3).

The global focus has for long been on the top 3 leading causes of neonatal mortality: prematurity, sepsis and perinatal asphyxia (4). Surgical interventions are important in improving the outcomes of many neonatal surgical diseases as timely interventions lead to better outcomes, improve quality of life and reduce disability. Successful management of neonates with surgical conditions requires a

multidisciplinary approach, especially in postoperative care. Factors such as sepsis control, care in the newborn intensive care unit (NICU) and availability of total parenteral nutrition (TPN) have been shown to improve the outcomes. Mortality rates from neonatal surgical cases remain high in most developing countries largely due to lack of NICU, scarcity of surgical specialists and unavailability of TPN (5–9).

High-income countries (HICs) continue to report lower mortality rates even with an apparent increase in neonatal surgical cases (2). Good surgical outcomes that have been achieved in HICs are feasible in developing countries only if some of the associated adverse factors are addressed (2, 3). A study done at Kenyatta National Hospital (KNH) in 2015 reported a mortality rate of 33% (10). This study set out to evaluate the outcomes (mortality and length of hospital

stay) of neonates requiring surgical interventions at Moi Teaching and Referral Hospital (MTRH).

## Methods

This was a prospective descriptive study carried out at the Newborn Unit of MTRH between February and November 2016. The study aimed to evaluate the outcomes (mortality and length of hospital stay) of neonates requiring surgical interventions. The study population comprised neonates with conditions requiring surgical interventions in the neonatal period. The unit has a capacity of 60 beds and provides care to neonates with medical and surgical problems. The approximate nurse to patient ratio is 1:20. The neonates are reviewed regularly by surgical specialists (general pediatric surgeons, neurosurgeons, orthopedic and ENT surgeons) upon admission. Once the decision for surgical interventions on their conditions was made, the neonates were recruited into the study consecutively, once written informed consent was obtained from the parents/guardians. Ethical approval was obtained from the Institutional Review and Ethics Committee. Data on maternal age, place and mode of delivery, newborn's age at admission, birth weight, surgical conditions, co-morbid conditions, treatment outcomes (discharge, death or referral to KNH for specialized care), surgical complications and time-to-initiation of oral feeds postoperatively were recorded. Antenatal history and laboratory parameters were also obtained. The length of hospital stay was determined following the final disposition of the newborn, which was either discharge after recovery, death or referral to KNH for specialized surgery. Data were analyzed using STATA® version 13 at 95% confidence interval. Data are presented using frequency tables. Simple logical regression tested for independent associations, and Mann-Whitney U test compared two medians. p-values of less than 0.05 were considered statistically significant.

## Results

During the study period 1,096 neonates were admitted to the unit: neonatal surgical conditions accounted for 11.3%. A total of 124 neonates were recruited into the study. The male to female ratio was 1.1:1. Most mothers attended antenatal clinics (82.3%), and nutritional supplements were issued to 84 (82.4%) mothers during pregnancy. None of the mothers received nutritional supplements during the first trimester. Routine antenatal ultrasound scans were done on only 4% of the mothers and no congenital anomaly was

detected in the ante-partum period. Most neonates (83%) were delivered in other health facilities and before they were referred to MTRH for surgical care; 9 (7%) were delivered at MTRH, and 12 (10%) at home. The median age at admission was 2 days (IQR 1, 5) with a range of 1 to 26 days; 59.7% of the neonates had birth weights of between 2.5 and 3.9 kg, 31.5% between 1.5 and 2.4 kg, and 4% were >3.9 kg.

### Surgical interventions offered

Most neonates (87.1%) who required surgical interventions had surgery done at MTRH. Of these 65(60.2%) were emergency surgical operations and 43(39.8%) were elective. Reasons cited for not performing surgery on some of the neonates ( $n=16$ ) in MTRH included: too sick to withstand surgery (43.8%), died before surgery (25%), were booked to be seen in pediatric surgical outpatient clinic (18.8%), and referred to KNH for specialized surgery (12.5%).

### Treatment outcomes

The overall mortality rate was 31.5%. Eighty-three (66.9%) neonates were discharged home upon recovery and 2(1.6%) were referred to KNH for specialized surgery.

### Spectrum of neonatal surgical conditions

Most neonatal surgical conditions (92%) were congenital anomalies: 70% had surgical conditions affecting the gastrointestinal system, 20% the central nervous system and 10% the musculoskeletal system. Table 1 shows the top 4 diagnoses in this study. The other 19(15.3%) neonates had diagnoses that included sacrococcygeal teratoma, pyomyositis, cystic hygroma, meconium peritonitis, brain abscess and submandibular abscess (Table 1).

Table 1: Main diagnoses, mortality and discharge/referral rates

Diagnosis	Frequency $n$ (%) $n=124$	Discharge / Referred to KNH $n$ (%) $n=85$	Mortality $n$ (%) $n=39$
Gastroschisis	33 (26.6)	14 (42.5)	19 (57.5)
Neural tube defects	25 (20.2)	23 (92)	2 (8)
ARM	25 (20.2)	17 (68)	8 (32)
Small bowel atresia	18 (14.5)	13 (72)	5 (28)
OA/TEF	4 (3.2)	1 (25)	3 (75)
Others	19 (15.3)	17 (89.5)	2 (10.5)

### Co-morbid conditions

Co-morbid conditions were reported in 94(75.6%) of neonates, with anemia being the leading co-morbid condition. Bivariate analysis showed a significant association between the presence of co-morbid conditions (respiratory distress,

electrolytes imbalance, acute kidney injury) and mortality,  $p=0.001$ , (Table 2).

Table 2: Bivariate analysis showing association between independent variables and mortality ( $n = 122$ )

	No = 83 (n/%)	Yes = 39 (n/%)	p-value
<b>ANC attendance</b>			
None	12 (54.5)	10 (45.5)	0.325
1–3 visits	59 (71)	24 (29)	
≥4 visits	12 (70.6)	5 (29.4)	
<b>Delivery place</b>			
Home	9 (75)	3 (25)	0.750*
Health facility	74 (67.3)	36 (32.7)	
<b>Birth weight</b>			
<2.5kg	20 (45.5)	24 (54.5)	0.001
≥2.5kg	63 (80.8)	15 (19.2)	
<b>Sepsis</b>			
Yes	33 (62.3)	20 (37.7)	0.230
No	50 (72.5)	19 (27.5)	
<b>Respiratory distress</b>			
Yes	12 (35.3)	22 (64.7)	0.001
No	71 (80.7)	17 (19.3)	
<b>Electrolyte imbalance</b>			
Yes	10 (33.3)	20 (66.7)	0.001
No	73 (79.3)	19 (20.7)	
<b>AKI</b>			
Yes	2 (25)	6 (75)	*0.013
No	81 (71)	33 (29)	

\*Fishers exact test

### **Surgical complications**

The surgical complication rate was 26.9%: wound sepsis (17.6%), post-operative hemorrhage (5.6%), burst abdomen (2.8%) and bowel perforations (1.9%).

### **Laboratory parameters**

Blood cultures isolated mainly *Klebsiella pneumoniae* species, and low platelet count was significantly associated with mortality ( $p<0.001$ ).

### **Length of hospital stay**

The median length of hospital stay was 14.5 days (IQR 6, 32) with a range of 0–102 days. Time-to-initiation of oral feeds postoperatively ranged from 1 to 24 days, and 51.3% of the neonates stayed nil per oral for more than 3 days postoperatively. Bivariate analysis using Mann-Whitney U test showed a significant association between time-to-initiation of oral feeds postoperatively and length of hospital stay ( $p<0.05$ ).

### **Factors associated with outcomes of neonates who required surgical interventions**

Antenatal clinic attendance was associated with decreased odds of mortality (OR 0.126; 95% CI 0.025–0.6429;  $p=0.013$ ) while neonatal sepsis, respiratory distress and electrolyte imbalance were associated with increased odds of mortality (OR 3.4, 95% CI 1.09–22.06,  $p=0.049$ ; OR 4.9, CI 0.91–11.61,  $p=0.001$ ; and OR 03.1, CI 1.21–31.60,  $p=0.029$ ; respectively (Table 3).

Table 3: Simple logistic regression showing association of independent variable with mortality

Variable	Odds ratio	p value	Confidence interval
Platelets ( $\times 10^9/L$ ) (50–150 vs <50)	0.2636	0.286	0.0227, 3.0598
Platelets ( $\times 10^9/L$ ) (>150 vs <50)	0.1199	0.042	0.0156, 0.9252
ANC (attendance vs no attendance)	0.1262	0.013	0.0248, 0.6429
Birth weight (kg) ( $\geq 2.5$ vs <2.5)	0.8463	0.792	0.2448, 2.9262
Sepsis (Yes vs No)	3.3573	0.049	0.9138, 11.6104
Respiratory distress (+ vs –)	4.9024	0.038	1.0896, 22.0572
Electrolyte imbalance (+ vs –)	6.1778	0.029	1.2077, 31.6020
AKI (+ vs –)	3.0720	0.381	0.2495, 37.8303

### **Discussion**

The median age at presentation to hospital was 2 days, which is late for most neonatal surgical conditions especially anterior abdominal wall defects and small bowel atresia. This could be a result of late referral by the peripheral health facilities. There could also have been delays in transport; MTRH has a large catchment population with some neonates coming from as far as Turkana and West Pokot counties, which have poor terrain.

Care of neonates with surgical conditions is limited to MTRH as the only tertiary facility in the region because pediatric surgeons who would offer such specialized care are few in the lower level hospitals. The median age at presentation in this study contrasts with that found in Nigeria which was 23.5 hours (11). The Nigerian study looked at neonates with anterior abdominal wall defects only, thus mothers could have rushed the neonates to hospital because of the dramatic presentation of those conditions.

The mortality rate in this study was 31.5%, which is almost similar to the rate in a study done at KNH whose overall mortality rate was 33% (10). KNH and MTRH are the only national referral hospitals in the country; the 2 centers are also tertiary level hospitals managing neonates with surgical

disorders referred from peripheral facilities within their catchment populations. The delays in presentation may have led to complications, more so, fluid and electrolyte imbalances, hence the higher mortality rate in both hospitals. The almost similar findings could also be due to the smaller number of neonates studied in both hospitals and the short durations of both studies.

A study done at MTRH in 2008 found an overall mortality rate of 24%, implying a shift in the outcomes over the 10 years. However, that study was retrospectively done over a 3-year period on a smaller sample of neonates with surgical conditions. It was carried out at the old unit which had a bed capacity of 20 compared with this study done at the new unit with a bed capacity of 60, and brings together different disciplines—neuro-surgery, orthopedics and plastic surgery (12). This increase in the number of admissions of neonates with surgical conditions over the years has led to congestion in the unit, which could explain the change in the mortality rate: neonatal sepsis is one of the factors significantly associated with mortality. This study reported a lower mortality rate than a study by Ugwu in Nigeria that reported a mortality rate of 48.2% after surgical interventions (5). Another study done by Kouame in Abidjan reported a higher mortality rate of 52% (13).

Lower mortality rates were reported in a survey to compare outcomes in low income countries (LICs) and high income countries (HICs). Mortality rates lower than 25% were reported in HICs, 50–75% in middle income countries (MICs) and more than 75% in some LICs (2). This discrepancy may be because the survey was carried out during an annual scientific conference with self-reported questionnaires from the respondents, thus the possibility exists of recall bias influencing the validity of the study findings. The lower mortality rates reported in HICs could also be because NICU and TPN are available; they are important adjuncts of the peri-operative care of surgical neonates.

The most common diagnosis was gastroschisis, in contrast to a study done in the same center 10 years earlier where ARM was the most common condition (34%) (12). It contrasts with the KNH study (10) which reported ARM at the top (19.2%) and next was anterior abdominal wall defects (17.2%). This could be explained by the global shift in the incidence of neonatal surgical conditions. Gastroschisis for instance is reported to have increased from 6% in 2003 to 15% in 2007 in Durban (6). The incidence of anterior abdominal wall defects has increased globally, with the incidence of gastroschisis increasing from 2.5/10,000 in 1994 to 4.4/10,000 in 2004, according to the British Register on anomalies (14). There was also a large catchment population for neonates with

gastroschisis in this study, with some coming from as far as Kiambu, Turkana and Nakuru counties, thus reflecting the rise in gastroschisis cases managed at MTRH.

Lack of proper use of nutritional supplements, such as folic acid, before conception and during pregnancy has also been associated with the occurrence of anterior abdominal wall defects and neural tube defects. Most neonates were delivered by mothers who did not take folic acid during the first trimester and in the preconceptional period as recommended. Over 50% of participants' mothers used supplements like folic acid and iron during the second trimester of pregnancy, when organogenesis is presumably already complete in the fetus, and thus use of these supplements could not have been of benefit in preventing neural tube defects.

Most participants were delivered in peripheral health facilities and later referred to MTRH for surgical care. This number compares with other studies as most of those studies were done in tertiary hospitals where such care of neonates with surgical conditions is offered (5, 7, 11, 12, 15, 16). Most referred neonates had poor outcomes with a mortality rate that was similar to other studies (5, 11, 12). Although ANC attendance was significantly associated with low mortality, attendance was inadequate with only 13.7% of mothers completing the recommended 4 ANC visits during pregnancy. Low ANC attendance in some studies and improper use of nutritional supplements in the periconception period as recommended could also have contributed to a rise in some conditions like gastroschisis and neural tube defects (5, 17, 18).

No neonatal surgical condition was diagnosed in utero, probably because almost half of the participants' mothers went for their first ANC visit in the second trimester and only 4% had an obstetric ultrasound done during pregnancy with no congenital anomaly being diagnosed prenatally. An explanation could be that most ultrasound scans at the peripheral health facilities are done by ultrasonographers who may not have the expertise to diagnose or identify some of the congenital anomalies.

Only 25% of participants had blood cultures done that established sepsis in 21% of the participants, and almost half the participants had presumptive diagnosis of probable sepsis. Most neonates who had complications following surgery had wound sepsis, a finding that could partly explain the prolonged hospital stay. This in turn could have led to hospital-acquired infection. Wound sepsis could also be due to a breach in the skin for some conditions like anterior abdominal defects and frequent handling due to daily cleaning and dressing of incision wounds. The presence of intravenous lines and naso-gastric tubes could

also predispose to neonatal sepsis. This finding compares with most studies where hospital-acquired sepsis was a leading cause of morbidity and mortality with poor outcomes associated with sepsis (5, 7, 8).

In this study most neonates (51.3%) stayed nil per oral for more than 3 days with a range of 1–24 days. These neonates would have benefitted from TPN, but it was not available in the unit. Neonates who had respiratory distress were four times more likely to die compared with those without respiratory distress, pointing towards the need to provide better respiratory support for them. Poor outcomes in most developing countries have been attributed to lack of supportive care such as the standard use of TPN and NICU services.

Of the neonates who did not have surgical intervention at MTRH, about a half of them died before surgery or were too sick to withstand surgery. These poor outcomes could be attributed to lack of NICU and proper ventilatory support. This is reflected in the finding of 25% of participants who died before surgery, and the significant number who were too sick to withstand surgery and who eventually died. Low platelet count, respiratory distress requiring CPAP or oxygen use, and electrolyte imbalance all related to poor outcomes. These findings compare with studies done in other centers that showed that sepsis, multiple organ failure leading to respiratory distress, and thrombocytopenia contributed most to mortality (17, 19, 20).

The median length of hospital stay was 14.5 days, which is in contrast to a study done in Nigeria with a median length of hospital stay of 10 days. Most participants in the Nigerian study had isolated anterior abdominal wall defects, mainly omphalocele major, that necessitated primary abdominal closure, hence the shorter hospital stay (11). This long hospital stay contrasts with a study done at KNH that had a median length of 10.2 days (10). The longer period of hospital stay in this study could be explained by the high number of neonates with gastroschisis who presented late and thus had to undergo staged closure, delaying recovery and prolonging hospital stay. The association between time of initiating oral feeds and length of hospital stay was also significant. This is evidenced by the 50% mortality rate among participants who stayed for more than 72 hours without enteral feeds, pointing to the need to initiate timely feeds postoperatively. The presence of co-morbid conditions like neonatal sepsis and neonatal jaundice could also have contributed to the longer hospital stay, as treatment of confirmed sepsis requires antimicrobial use for a minimum of 10–14 days. Postoperative complications like wound

sepsis and burst abdomen may also have necessitated re-operation, which may further prolong the duration of hospital stay.

## Conclusions

The overall mortality rate of neonates requiring surgical interventions at MTRH was 31.5%, and their median length of hospital stay was 14.5 days. Neonatal sepsis, respiratory distress and electrolyte imbalance were significantly associated with increased odds of mortality.

## Acknowledgements

We acknowledge the contributions of Dr Ann Mwangi in data analysis and management.

## References

1. WHO | Infant, Newborn, [www.int.topics/infant\\_newborn/en/](http://www.int.topics/infant_newborn/en/)
2. Wright NJ, Zani A, Ade-Ajayi N. Epidemiology, management and outcome of gastroschisis in Sub-Saharan Africa: Results of an international survey. *Afr J Paediatr Surg.* 2015; 12(1):1–6.
3. Chirdan LB, Ameh EA. Untreated surgical conditions: Time for global action. *Lancet.* 2012; 380(9847):1040–41.
4. World Health Organization. The Global Burden of Disease: 2004 Update. Geneva: WHO Press; [update.www.int/healthinfo/global\\_burden\\_disease/2004\\_report](http://update.www.int/healthinfo/global_burden_disease/2004_report)
5. Ugwu RO, Okoro PE. Pattern, outcome and challenges of neonatal surgical cases in a tertiary teaching hospital. *Afr J Paediatr Surg.* 2013 (3):226–30.
6. Sekabira J, Hadley GP. Gastroschisis: A Third World perspective. *Pediatr Surg Int.* 2009; 25:327–329.
7. Shakya VC, Agrawal CS, Shrestha P, et al. Management of *Jejunal atresia: An Experience at Eastern Nepal.* BMC Surg. 2010.
8. Ameh EA, Chirdan LB. Neonatal intestinal obstruction in Zaria, Nigeria. *East Afr Med J.* 2000.
9. Taguchi T, Nagata K, Kinoshita Y, et al. Progress in and outcomes of neonatal surgery Over the Past 50years. *Nihon Geka Gakkai Zasshi.* 2014; 115(6):306–11.
10. Ajanja SO. Pattern of presentation of neonatal surgical disease and outcomes of surgery at Kenyatta National Hospital. MMed thesis, University of Nairobi. Available at [www.uonbi.ac.ke](http://www.uonbi.ac.ke).
11. Abdur-Rahman LO, Abdulrasheed NA, Adeniran JO. Challenges and outcomes of management of anterior abdominal wall defects in a Nigerian tertiary hospital. *Afr J Paediatr Surg.* 2011 (2):159–63.
12. Tenge-Kuremu R, Kituyi PW, Tenge CN, et al. Neonatal surgical emergencies at Moi Teaching and Referral Hospital in Eldoret, Kenya. *ECAMJ.* 2006; 12(2):36–39.
13. Kouame BD, N'guetta-Brou IA, Kouame GS, et al. Epidemiology of congenital abnormalities in West Africa: Results of a descriptive study in a teaching hospital in Abidjan, Cote d'Ivoire. *Afr J Paediatr Surg.* 2015; (1):51–5.
14. British Isles Network of Congenital Anomaly Registers. <https://www.gov.uk/.../the-national-congenital-anomaly-and-rare-disease-registration>. 2015.

15. Mouafo Tambo FF, Chiabi A, Ngowe Ngowe M, et al. Mortality of neonatal surgical emergencies at the Gynecology, Obstetric and Pediatric Hospital of Yaoundé, Cameroon. *Med Trop (Mars)*. 2011; 71(2):206–7.
16. Ouédraogo I, Kaboré R, Napon Madina A, et al. Epidemiology of neonatal surgical emergencies in Ouagadougou. *Arch Pediatr*. 2015; (2):130–4.
17. Tarcă E, Aprodu S. Past and present in omphalocele treatment in Romania. *Chirurgia (Bucur)*. 2014; 109(4):507–13.
18. Botto LD, Mulinare J, Erickson JD. Occurrence of omphalocele in relation to maternal multivitamin use: A population-based study. *Pediatrics*. 2002; 109(5):40–9.
19. Arnold M, Moore SW, Sidler D, et al. Long-term outcome of surgically managed necrotizing enterocolitis in a developing country. *Pediatr Surg Int*. 2010; 26(4):355–60.
20. Kuradusenge P, Kuremu RT, Jumbi G, et al. Pattern of anorectal malformation and early outcomes of management at Moi Teaching and Referral Hospital, Eldoret, Kenya. *EAMJ*, 2004; 91(12):430–34.